(19) World Intellectual Property Organization International Bureau



(43) International Publication Date 28 December 2000 (28.12.2000)

PCT

(10) International Publication Number WO 00/78715 A1

TOMINAGA, Mineko [BR/BR]; CEP-São Paulo,

SP (BR). DE MATTOS PAIVA, Antonio, Cechelli

[BR/BR]; CEP-São Paulo, SP (BR), DOS REIS BAR-BOSA, Simone [BR/BR]; CEP-São Paulo, SP (BR).

MARCHETTO, Reinaldo [BR/BR]; CEP-São Paulo, SP (BR), SCHREIER, Shirley [BR/BR]; CEP-São Paulo, SP

EURY PEREIRA, Luna-Filho; Avenida

- (51) International Patent Classification7: C07D 207/46, 295/24
- (21) International Application Number: PCT/BR00/00068
- (22) International Filing Date: 21 June 2000 (21.06.2000)
- (25) Filing Language:

English English

(26) Publication Languag

CEP-70741-901 Brasilia, DF (BR).

- (30) Priority Data: PI 9903137-0 1/ 22 June 1999 (22.06.1999)
- (71) Applicant (for all designated States except US): CON-SELHO NACIONAL DE DESENVOLVIMENTO
- W3 Norte, Quadra 507/B, Lote 2, SEPN, Suite 211, CEP-70741-901 Brasilia, DF (BR). (81) Designated States (national): JP, US.

 - (84) Designated States (regional): European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).

Published:

(74) Agent:

- With international search report.
- Before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments.


[Continued on next page]

(72) Inventors: and

(75) Inventors/Applicants (for US only); NAKAIE, Clevis, Ryuichi [BR/BR]; CEP-São Paulo, SP (BR).

CIENTÍFICO E TECNOLÓGICO - CNPO [BR/BR]: Avenida W3 Norte, Quadra 507/B, Lote 2, SEPN,

(54) Title: SYNTHESIS OF A NOVEL PARAMAGNETIC AMINO ACID DERIVATIVE (EPM-5) FOR LABELLING CHEMI-CAL AND BIOLOGICAL MACROMOLECULES



Structure of Fmoc-Poac.

8

Com Sam of

(57) Abstract: The present invention refers to the synthesis and application of 2,2,5,5-tetramethylpyrrolidine-N-oxyl-(9-fluorenylmethyloxycarbonyl)-3-amine-4-carboxylic acid, a novel paramagnetic (spin label) amino acid derivative denominated as Fmoc-POAC whose structure is seen in Figure 1. Fmoc-POAC can be coupled to peptide sequences and other systems. It can be inserted anywhere in a peptide segment, even at an internal position, after removal of its temporary amine protecting group, Fmoc. Owing to its pyrrolidine structure, this molecule may induce differentiated conformations if compared with the normal alpha-amino acids, thus becoming a valuable probe for stretural-biological activity of certain peptides. The POAC-angiotensin II analogue was synthesized as a model according to the use of the chemical derivative.